0.5 bet

<div class="hwc kCrYT" style="padding-bottom:12px;padding-top:
0px"><div><div><div><div><div><div>&
lt;div>How often should I get the O-Shot Treatment? The O-Shot can be a <s
pan>one-time treatment for up to two years of improved sexual we
llness. However, studies suggest that a second O-Shot six to twelve months after
 the first can extend the results achieved.</div></div></div>&
lt;/div></div><div></div><div><a data-ved="2a
hUKEwjrveaj_s-DAxWqMUQIHRWqAyIQFnoECAEQBg" href="{href}"><s
pan><div>O-Shot Treatment In Houston, TX | Northwest Women&
#39;s Center</div><div>nwwomen
s : o-shotin-houston-tx</div></div></div&
gt;</div><div><div><div><a data-ved=&quot
;2ahUKEwjrveaj_s-DAxWqMUQIHRWqAyIQzmd6BAgBEAc" href="{href}">0
.5 bet</div></div></div></div>&lt
;div class="hwc kCrYT" style="padding-bottom:12px;padding-top:0px
"><div><div><div><div></div>&l
t;h2><div>What Are the Adverse Effects of O-shot?</span&
gt;</div></h2></div><div></div><di
v><div><div><div><div>Urgent or more freq
uent urination.</div></div></div></div><
/div><div></div><div><div><div><div><s
pan><div>Pinching, pressure or pain during injection (Topical numbing a) Tj T*□□BT /F1 12 Tf

gt;</div></div><div></div><div><div><div&
gt;<div><div>Spotting.</div></div&
gt;</div></div></div><div></div><div><div
><div><div><div>A feeling of vaginal "full
ness"</div></div></div></div></div&
gt;<div><div><a data-ved="2ahUKEwjrv
eaj_s-DAxWqMUQIHRWqAyIQFnoECAEQDQ" href="{href}">&
lt;span>O-Shot Safety - Coyle Institute&l
t;/span>□<a data-ved="2ahUKEwjrveaj_s-DAxWqMUQIHRWqAyIQl
qUEegQIARAO" href="{href}">coyleinstit
ute : o-shot-safety</div&gt
;</div></div></div><div><div><div&
gt;<span><a data-ved="2ahUKEwjrveaj_s-DAxWqMUQIHRWqAyIQzmd6BAgBEA8